

EDITORIALS



Improving outcomes from in-hospital cardiac arrest

Make sure local practice is informed by the latest guidelines

Keith Couper *postdoctoral research fellow*^{1 2}, Gavin D Perkins *professor*^{1 2}

¹Warwick Medical School, University of Warwick, Coventry CV4 7AL, UK; ²Heart of England NHS Foundation Trust, Birmingham, UK

Over 200 000 adults a year sustain a cardiac arrest while in hospital in the United States.¹ Most trials have taken place outside hospital,² yet the etiology, patient characteristics, time to treatment, and outcomes are quite different from cardiac arrests occurring in inpatients. Clinical guidelines for in-hospital resuscitation are therefore mainly drawn from the extrapolation of findings from out-of-hospital trials, observational studies, and consensus of expert opinion coordinated through the International Liaison Committee for Resuscitation.³

Given the cost, logistical, and ethical challenges of conducting randomized trials in cardiac arrest, the use of high quality observational data to provide insights into the effectiveness of treatments is attractive. The main limitation of observational studies is the risk that the outcome is affected by both the treatment allocation and other factors that influence the treatment allocation. Propensity scoring methods have been growing in popularity as a way of reducing confounding related to measured variables.

In critically ill patients, well conducted propensity score analyses generally agree with findings from randomized controlled trials, although the effect size may vary.⁴ A key limitation nevertheless remains the bias caused by unmeasured confounders. This was illustrated in two propensity analyses using data from the same registry on the effect of adrenaline on survival from out-of-hospital cardiac arrest. The studies yielded diametrically opposing results through small differences in the variables included in the propensity scoring model.⁵

In *The BMJ*, two linked research papers^{6 7} use propensity score analyses of data from the American Heart Association's (AHA) Get With The Guidelines-Resuscitation (GWTG-R) registry to examine the association between different treatment strategies for cardiac arrest and patient outcomes in in-hospital patients with shock refractory ventricular fibrillation (VF) or pulseless ventricular tachycardia (VT).

Bradley and colleagues⁶ explored differences in outcomes between defibrillator shocks delivered in rapid succession (stacked shocks), which formed part of the AHA guidelines in 2000, and the strategy of delivering a single shock strategy followed by 2 minutes of chest compressions, which was introduced in 2005 (fig 1).⁸ Their analysis showed slow adoption

of the single shock sequence (increase from 30% in 2006 to 60% in 2012). Although unadjusted analyses showed better hospital survival with stacked shocks, there was no difference after adjustment for propensity scoring (adjusted risk ratio 0.89, 95% confidence interval 0.78 to 1.01).




Fig 1 Difference between defibrillator shock strategies (stacked shocks v single shock sequence), as recommended in the AHA guidelines before and after 2005

The change in shock sequence from stacked to single shocks was introduced concurrently with the recommendation that adrenaline (epinephrine) administration was deferred until after 2 minutes of cardiopulmonary resuscitation (CPR) after the time of rhythm reanalysis and close to delivery of a second shock (fig 1). The premise for deferring adrenaline was to avoid giving it blindly before determining the response to initial defibrillation and potentially precipitating refrillation.

Andersen and colleagues⁷ explored the association between early (within 2 minutes) and deferred (or no) adrenaline in patients enrolled in the registry from 2006 (to avoid contamination with changes in shock sequences). In contrast to previous work from the GWTG-R group where early adrenaline seemed to improve outcomes in patients with non-shockable rhythms,⁹ the early administration of adrenaline to patients with shock refractory VF or VT was associated with reduced survival to hospital discharge (adjusted odds ratio 0.70, 95% confidence interval 0.59 to 0.82).

The key strengths of both studies are the use of the high quality GWTG-R registry, which collates data from over 300 hospitals in the USA and serves as a rich source of high quality information on treatments and outcomes from in-hospital cardiac

arrest.¹⁰ While the breadth of coverage across the USA is a major strength, its primary purpose as a quality improvement registry necessarily limits the depth and specificity of information that is available for analysis. In both studies, researchers were reliant on using timings to draw inferences on the treatments being administered rather than extracting specific information about single versus stacked shocks or about timing of adrenaline in relation to shock sequence. Without such granular information, the slow adoption of deferred shock strategy identified by Bradley and colleagues and its interaction with the timing of adrenaline and other unmeasured and potentially confounding variables (eg, changes in ratio of compressions to ventilations, improving CPR quality) make it difficult to conclude a causal association between the studied interventions and survival.

So should the findings from these studies alter your practice? Yes, the finding of widespread non-adherence with clinical guidelines should prompt those responsible for organizing or delivering advanced life support to review their practice and ensure that it is informed by the latest clinical guidelines. While the jury remains out on the overall safety or effectiveness of adrenaline in cardiac arrest,¹¹ these data suggest that if adrenaline is given, in accordance with current guidelines, it should be deferred until at least after the second shock has been delivered.

Competing interests: We have read and understood the BMJ policy on declaration of interests and declare the following interests: KC is supported by a National Institute for Health Research (NIHR) postdoctoral fellowship; GDP is supported as a NIHR senior investigator, and is co-chair of the International Liaison Committee for Resuscitation. Provenance and peer review: Commissioned, not externally peer reviewed.

- 1 Mozaffarian D, Benjamin EJ, Go AS, et al. American Heart Association Statistics Committee and Stroke Statistics Subcommittee. Heart Disease and Stroke Statistics-2016 Update: A Report From the American Heart Association. *Circulation* 2016;133:e38-360. doi:10.1161/CIR.0000000000000366 pmid:26673558.
- 2 Whitehead L, Perkins GD, Clarey A, Haywood KL. A systematic review of the outcomes reported in cardiac arrest clinical trials: the need for a core outcome set. *Resuscitation* 2015;88:150-7. pmid:25497393.
- 3 Nolan JP, Hazinski MF, Aickin R, et al. Part 1: Executive summary: 2015 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science with Treatment Recommendations. *Resuscitation* 2015;95:e1-31. doi:10.1016/j.resuscitation.2015.07.039 pmid:26477703.
- 4 Kitsios GD, Dahabreh IJ, Callahan S, Paulus JK, Campagna AC, Dargin JM. Can we trust observational studies using propensity scores in the critical care literature? A systematic comparison with randomized clinical trials. *Crit Care Med* 2015;43:1870-9. doi:10.1097/CCM.0000000000001135 pmid:26086943.
- 5 Olasveengen TM. Adrenaline for out of hospital cardiac arrest? *BMJ* 2013;347:f7268. doi: 10.1136/bmj.f7268 pmid:24326888.
- 6 Bradley SM, Liu W, Chan PS, et al. American Heart Association's Get With The Guidelines-Resuscitation Investigators. Defibrillation time intervals and outcomes of cardiac arrest in hospital: retrospective cohort study from Get With The Guidelines-Resuscitation registry. *BMJ* 2016;353:i1653.
- 7 Andersen LW, Kurth T, Chase M, et al. American Heart Association's Get With The Guidelines-Resuscitation Investigators. Early administration of epinephrine (adrenaline) in patients with cardiac arrest with initial shockable rhythm in hospital: propensity score matched analysis. *BMJ* 2016;353:i1577.
- 8 Hazinski MF, Nadkarni VM, Hickey RW, O'Connor R, Becker LB, Zaritsky A. Major changes in the 2005 AHA Guidelines for CPR and ECC: reaching the tipping point for change. *Circulation* 2005;112(Suppl):IV206-11. doi:10.1161/CIRCULATIONAHA.105.170809 pmid: 16314349.
- 9 Donnino MW, Saliccioli JD, Howell MD, et al. American Heart Association's Get With The Guidelines-Resuscitation Investigators. Time to administration of epinephrine and outcome after in-hospital cardiac arrest with non-shockable rhythms: retrospective analysis of large in-hospital data registry. *BMJ* 2014;348:g3028. doi:10.1136/bmj.g3028 pmid: 24846323.
- 10 Peberdy MA, Kaye W, Ornato JP, et al. Cardiopulmonary resuscitation of adults in the hospital: a report of 14720 cardiac arrests from the National Registry of Cardiopulmonary Resuscitation. *Resuscitation* 2003;58:297-308. doi:10.1016/S0300-9572(03)00215-6 pmid: 12969608.
- 11 Perkins GD, Cottrell P, Gates S. Is adrenaline safe and effective as a treatment for out of hospital cardiac arrest? *BMJ* 2014;348:g2435. doi:10.1136/bmj.g2435 pmid:24709574.

Published by the BMJ Publishing Group Limited. For permission to use (where not already granted under a licence) please go to <http://group.bmj.com/group/rights-licensing/permissions>